



Thank you for selecting our infusion services team to care for your patient. Please provide ALL information listed below to ensure that we can process orders and schedule your patient for treatment without delay.

Part A- Patient scheduling and contact information:

Patient Name (Last, First): _____ Date of Birth: _____

Patient Contact Information and Phone Number (s): _____

Ordering Provider Name (Print): _____

Provider Clinic or Service Address: _____

Clinic or Service Phone Number: _____ Clinic or Service Fax Number: _____

Diagnosis (include ICD 10 codes): _____

Medication and Service Requested- list J-Code/ CPT code if known: _____

Date Service is Requested to Begin: _____ **Date Service is Expected to End:** _____

Order will expire 1 year from date of provider signature unless "date service is expected to end" is earlier.

Part B- Insurance and Prior Authorization. Any non-PeaceHealth provider must obtain prior authorization prior to service. Attach a copy of authorization documentation received from insurance payer when submitting orders.

Insurance (Payer) Company: _____

Prior Authorization Number and Conditions: _____

Prior Authorization Expiration Date: _____

Insurance (Payer) Contact Phone Number: _____

Part C- Elements needed to guide medication therapy are included with request for service:

- ☐ Orders and instruction (use the PeaceHealth approved ordering form if you are not a PeaceHealth provider) are complete and include provider signature at the bottom of each page. Check the boxes of ALL orders you would like to activate.
- ☐ For blood products, PeaceHealth Blood and Transfusion Consent form is signed and dated by the provider and the patient.

If information is located outside of PeaceHealth's electronic medical record system attach the following:

- ☐ A list of current medications reconciled by patient provider is available and includes a list of known allergies.
- ☐ Recent progress notes from ordering provider.
- ☐ A copy of relevant laboratory results and other appropriate supporting documentation.

IMPORTANT MESSAGE TO PROVIDERS: To reduce delays in treatment and added phone calls, you may participate to utilize PeaceHealth preferred medication formulary options by signing this document. A clinical pharmacist will adjust orders to align with PeaceHealth medical staff approved medication formulary options, policies, and procedures.

I agree to utilize PeaceHealth preferred medication formulary options, policies and procedures that have been authorized by PeaceHealth Medical Staff. This agreement will be issued for the duration of active orders contained within this treatment plan.

PROVIDER SIGNATURE: _____ **DATE:** _____ **TIME:** _____

FAX completed service request and orders to: PHMC OP Infusion and Nursing Services 541-902-1649



Abatacept (Orencia) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content
For Admission to Service	Provider Instruction – Review information below and address requirements for admission to service: - Provider has screened patient for history of tuberculosis and hepatitis B infection within 2 years of starting Abatacept (Orencia) therapy: • Date of hepatitis screening: _____ • Date of tuberculosis screening: _____
Supportive Care	☒ Abatacept (Orencia) IV infusion in NS 100 ml infused over 30 minutes: Select Dose: ☐ 500 mg (recommended for weight less than 60 kg) ☐ 750 mg (recommended for weight 60-100 kg) ☐ 1000 mg (recommended for weight greater than 100 kg) Select Frequency: ☐ Initiation regimen at 0, 2 and 4 weeks followed by maintenance infusion every 4 weeks ☐ Maintenance infusion every 4 weeks ☐ Maintenance infusion every _____ weeks (indicate frequency) Additional order instruction: ☒ Use 0.2-to-1.2-micron low protein-binding in-line filter.
Labs	☐ CBC with automated differential once prior to beginning treatment and every _____ weeks ☐ Comprehensive metabolic panel once prior to beginning treatment and every _____ weeks ☒ Treatment lab instructions – Provider approves to release and draw labs 2 days pre and post this planned treatment date.
Nursing Orders	☒ Confirm patient has negative PPD/tuberculosis and hepatitis B screenings prior to starting treatment. ☒ FOR INFUSION/ALLERGIC REACTION: Slow or stop abatacept infusion. Flush abatacept line with normal saline before administering PRN medications. Vital signs: every 15-30 minutes as needed. If reaction resolves, then resume infusion at half the previous rate. If reaction worsens discontinue infusion, maintain IV site until vital signs and conditions become stable, notify MD. ☒ Anaphylaxis (wheezing, dyspnea, hypotension, angioedema, chest pain, tongue swelling) and patient does not respond to diphenhydramine and/or methylprednisolone; give epinephrine, notify MD, and transfer to ED as needed. ☒ Patient may be discharged when vital signs are stable, patient does not display any evidence of adverse reaction, and infusion is complete or discontinued.
Nursing IV Access and Maintenance	Select the most appropriate option below: ☒ Insert PERIPHERAL IV as needed and flush (unless provider selects option for a central line). ☒ Sodium chloride 0.9% (NS) flush 10 mL IV once as needed for line care. ☐ Access and use NON-PICC Central Line/CVAD ☒ Initiate Central Line (non-PICC) maintenance protocol. ☒ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care, before and after medication administration, at discharge, and at de-access (sterile NS for Port-a-Cath access) ☒ Sodium chloride 0.9% (NS) flush 20 mL IV as needed for line care post lab draw. ☒ Heparin, porcine (PF) 100 unit/mL flush 5 mL IV as needed for line care, for de-access.

Practitioner Signature: _____ **Date of Order:** _____ **Time:** _____

Final page of orders must include signature of the ordering practitioner, date, and time.



Abatacept (Orencia) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content
	☒ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for occluded central line catheters- For clearing central line catheter. Reconstitute with 2.2 mL Sterile Water to the vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Instill medication in non-functional lumen. Do not use lumen while dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time not to exceed 120 min. Use second dose of Alteplase (Cathflo) if catheter not patent after 120 min. If the catheter is functional, aspirate and waste the medication and residual clot prior to flushing the line. ☐ Access and use <u>PICC</u> Central Line/CVAD ☒ Initiate PICC maintenance protocol. ☒ Change PICC line dressing weekly and as needed. ☒ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care, and before and after medication administration. ☒ Sodium chloride 0.9% (NS) flush 20 mL IV as needed for line care post lab draw ☒ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for occluded central line catheters- For clearing central line catheter. Reconstitute with 2.2 mL Sterile Water to the vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Instill medication in non-functional lumen. Do not use lumen while dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time not to exceed 120 min. Use second dose of Alteplase (Cathflo) if catheter not patent after 120 min. If the catheter is functional, aspirate and waste the medication and residual clot prior to flushing the line.
As Needed Medications	Standard As Needed Medications: ☒ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care. ☒ Sodium chloride 0.9% 500 mL continuous infusion at 25 mL/hour IV as needed for therapy administration (i.e., blood products, chemotherapy, potassium administration).
Emergency Medications	If patient has symptoms of anaphylaxis (wheezing, dyspnea, hypotension, angioedema, chest pain, or tongue swelling), discontinue infusion and initiate standard emergency response procedures. ☒ Standard Emergency Medications: ☒ Diphenhydramine (Benadryl) injection 25-50 mg IV once as needed for mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (greater than or equal to 20 points in SBP), nausea, urticaria, chills, pruritic). - Administer 50 mg IV if patient has NOT had diphenhydramine within 2 hours of reaction - Administer 25 mg IV if patient has had diphenhydramine within 2 hours of reaction, if reaction doesn't resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and notify provider. ☐ Albuterol 90 mcg/actuation inhaler 2 puffs once as needed for wheezing, shortness of breath associated with infusion reaction and contact provider. Administer with a spacer if available.

Practitioner Signature: _____ **Date of Order:** _____ **Time:** _____

Final page of orders must include signature of the ordering practitioner, date, and time.



Abatacept (Orencia) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content
	☒ MethylPREDNISolone (Solu-Medrol) injection 125 mg IV once as needed for shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (greater than or equal to 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist after administration of diphenhydramine (Benadryl) and notify provider. ☒ EPINEPHrine (Adrenalin) injection 0.5 mg IM once as needed for severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (greater than or equal to 40 points in SBP), shortness of breath with wheezing and O2 Sat less than 90%) and notify provider.
Referral	☒ Ambulatory referral to OP Infusion Services
PHMC Outpatient Infusion Contact Information	PROVIDER – PLEASE SIGN, DATE AND TIME ORDERS AND RETURN TO: PeaceHealth Peace Harbor Medical Center Outpatient Infusion Services Department 400 Ninth Street Florence, OR 97439 Contact Phone: 541-902-6019 and FAX 541-902-1649
Authorization by Verbal or Telephone Order	Person giving verbal or telephone order: _____ Person receiving verbal or telephone order: _____ ☐ Check to indicate verbal or telephone orders have been read back to confirm accuracy

Practitioner Signature: _____ Date of Order: _____ Time: _____

Final page of orders must include signature of the ordering practitioner, date, and time.