



Thank you for selecting our infusion services team to care for your patient. Please provide ALL information listed below to ensure that we can process orders and schedule your patient for treatment without delay.

Part A- Patient scheduling and contact information:

Patient Name (Last, First): _____ Date of Birth: _____

Patient Contact Information and Phone Number(s): _____

Ordering Provider Name (Print): _____

Provider Clinic or Service Address: _____

Clinic or Service Phone Number: _____ Clinic or Service Fax Number: _____

Diagnosis (include ICD 10 codes): _____

Medication and Service Requested - list J-Code/CPT code if known: _____

Date Service is Requested to Begin: _____ **Date Service is Expected to End:** _____

Order will expire 1 year from date of provider signature unless "date service is expected to end" is earlier.

Part B- Insurance and Prior Authorization. Any non-PeaceHealth provider must obtain prior authorization prior to service. Attach a copy of authorization documentation received from insurance payer when submitting orders.

Insurance (Payer) Company: _____

Prior Authorization Number and Conditions: _____

Prior Authorization Expiration Date: _____

Insurance (Payer) Contact Phone Number: _____

Part C- Elements needed to guide medication therapy are included with request for service:

- ☐ Orders and instruction (use the PeaceHealth approved ordering form if you are not a PeaceHealth provider) are complete and include provider signature at the bottom of each page. Check the boxes of ALL orders you would like to activate.
- ☐ For blood products, PeaceHealth Blood and Transfusion Consent form is signed and dated by the provider and the patient.

If information is located outside of PeaceHealth's electronic medical record system attach the following:

- ☐ A list of current medications reconciled by patient provider is available and includes a list of known allergies.
- ☐ Recent progress notes from ordering provider.
- ☐ A copy of relevant laboratory results and other appropriate supporting documentation.

IMPORTANT MESSAGE TO PROVIDERS: To reduce delays in treatment and added phone calls, you may participate to utilize PeaceHealth preferred medication formulary options by signing this document. A clinical pharmacist will adjust orders to align with PeaceHealth medical staff approved medication formulary options, policies, and procedures.

I agree to utilize PeaceHealth preferred medication formulary options, policies and procedures that have been authorized by PeaceHealth Medical Staff. This agreement will be issued for the duration of active orders contained within this treatment plan.

PROVIDER SIGNATURE: _____ **DATE:** _____ **TIME:** _____

FAX completed service request and orders to: PHMC OP Infusion and Nursing Services 541-902-1649



DAPTOmycin (Cubicin) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content																																																															
For Admission to Service	Provider Instruction – Daptomycin is a restricted antimicrobial. Approved indications include: - Serious infections due to MRSA, Coagulase negative Staphylococcus (CoNS), vancomycin-resistant organisms (VRE) including bacteremia, endocarditis, osteomyelitis, septic arthritis, prosthetic joint infections, cardiac device infections. Do not use for pneumonia (drug is inactivated by surfactant). - Continuation of therapy from outside hospital or outpatient use and prescribed by an ID clinician Reason for ordering (required): ☐ ID Consult ☐ Other reason: _____																																																															
Labs	☒ CBC with automated differential once prior to beginning treatment and weekly ☒ CMP once prior to beginning treatment and weekly ☒ CPK once prior to beginning treatment and weekly. Pharmacist can increase monitoring as needed up to three times weekly in at risk patients (i.e., HMG-CoA reductase inhibitor treatment, renal impairment). ☐ ESR once prior to beginning treatment and weekly ☐ CRP once prior to beginning treatment and weekly ☒ Provider approves to release and draw labs 2 days pre and post this planned treatment date.																																																															
Supportive Care	☒ DAPTOmycin (Cubicin) IV infusion every 24 hours over 30 minutes. Select Dose: ☐ 6 mg/kg ☐ 8 mg/kg ☐ 10 mg/kg ☒ Pharmacist may adjust dose and dosing interval based on the following table and renal function: <table border="1"> <thead> <tr> <th>Creatinine Clearance</th><th colspan="6">Indication</th></tr> </thead> <tbody> <tr> <td></td><td colspan="2"> - Skin/Soft Tissue - Surgical prophylaxis (x 1 dose) </td><td colspan="2"> - Uncomplicated MRSA/CoNS bacteremia - Osteomyelitis </td><td colspan="2"> - Enterococcal bacteremia - Persistent bacteremia - Infective endocarditis - CNS </td></tr> <tr> <td>Greater than or equal to 30 mL/min</td><td colspan="2">6 mg/kg q24h</td><td colspan="2">8 mg/kg q24h</td><td colspan="2">10 mg/kg q24h</td></tr> <tr> <td>Less than 30 mL/min</td><td colspan="2">6 mg/kg q48h</td><td colspan="2">8 mg/kg q48h</td><td colspan="2">10 mg/kg q48h</td></tr> <tr> <td></td><td>Dosing Weight</td><td>Dose</td><td>Dosing Weight</td><td>Dose</td><td>Dosing Weight</td><td>Dose</td></tr> <tr> <td></td><td>40-50 kg</td><td>250 mg</td><td>40-45 kg</td><td>250 mg</td><td></td><td></td></tr> <tr> <td></td><td>51-100 kg</td><td>500 mg</td><td>41-75 kg</td><td>500 mg</td><td>40-60 kg</td><td>500 mg</td></tr> <tr> <td></td><td>101-145 kg</td><td>750 mg</td><td>76-105 kg</td><td>750 mg</td><td>61-85 kg</td><td>750 mg</td></tr> <tr> <td></td><td>greater than 145 kg</td><td>1000 mg</td><td>greater than 105 kg</td><td>1000 mg</td><td>greater than 84 kg</td><td>1000 mg</td></tr> </tbody> </table>	Creatinine Clearance	Indication							- Skin/Soft Tissue - Surgical prophylaxis (x 1 dose)		- Uncomplicated MRSA/CoNS bacteremia - Osteomyelitis		- Enterococcal bacteremia - Persistent bacteremia - Infective endocarditis - CNS		Greater than or equal to 30 mL/min	6 mg/kg q24h		8 mg/kg q24h		10 mg/kg q24h		Less than 30 mL/min	6 mg/kg q48h		8 mg/kg q48h		10 mg/kg q48h			Dosing Weight	Dose	Dosing Weight	Dose	Dosing Weight	Dose		40-50 kg	250 mg	40-45 kg	250 mg				51-100 kg	500 mg	41-75 kg	500 mg	40-60 kg	500 mg		101-145 kg	750 mg	76-105 kg	750 mg	61-85 kg	750 mg		greater than 145 kg	1000 mg	greater than 105 kg	1000 mg	greater than 84 kg	1000 mg
Creatinine Clearance	Indication																																																															
	- Skin/Soft Tissue - Surgical prophylaxis (x 1 dose)		- Uncomplicated MRSA/CoNS bacteremia - Osteomyelitis		- Enterococcal bacteremia - Persistent bacteremia - Infective endocarditis - CNS																																																											
Greater than or equal to 30 mL/min	6 mg/kg q24h		8 mg/kg q24h		10 mg/kg q24h																																																											
Less than 30 mL/min	6 mg/kg q48h		8 mg/kg q48h		10 mg/kg q48h																																																											
	Dosing Weight	Dose	Dosing Weight	Dose	Dosing Weight	Dose																																																										
	40-50 kg	250 mg	40-45 kg	250 mg																																																												
	51-100 kg	500 mg	41-75 kg	500 mg	40-60 kg	500 mg																																																										
	101-145 kg	750 mg	76-105 kg	750 mg	61-85 kg	750 mg																																																										
	greater than 145 kg	1000 mg	greater than 105 kg	1000 mg	greater than 84 kg	1000 mg																																																										

Practitioner Signature: _____ **Date of Order:** _____ **Time:** _____

Final page of orders must include signature of the ordering practitioner, date, and time.



DAPTOmycin (Cubicin) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content
Nursing Orders	☒ RN to notify provider if CPK > 1000 units/L with unexplained signs and symptoms of myopathy and/or if CPK > 2000 units/L. ☒ At the end of treatment, contact provider to address removal of PICC line.
Nursing IV Access and Maintenance	Select the most appropriate option below: ☒ Insert PERIPHERAL IV as needed ☒ Sodium chloride 0.9% (NS) flush 10 mL IV once as needed for line care ☐ Access and use NON-PICC Central Line/CVAD as needed and confirm patency ☒ Initiate Central Line (Non-PICC) maintenance protocol ☒ Sodium chloride 0.9% (NS) injection 10 mL IV as needed for line care before and after medication administration, at discharge, and at de-access ☒ Sodium chloride 0.9% (NS) injection 20 mL IV once as needed for line care post lab draw ☒ Heparin, porcine (PF) 100 unit/mL flush 5 mL IV as needed for line care, for de-access ☒ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for clearing central line catheter. Reconstitute with 2.2 mL sterile water for injection to the vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1mg/mL. Instill medication in non-functional lumen. Do not use lumen while dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time not to exceed 120 minutes. Use second dose of Alteplase (Cathflo) if catheter not patent after 120 minutes. If the catheter is functional, aspirate and waste the medication and residual clot prior to flushing the line. ☐ Access and use PICC Central Line/CVAD as needed and confirm patency ☒ Initiate PICC maintenance protocol ☒ Change PICC line dressing weekly and as needed ☒ Sodium chloride 0.9% (NS) injection 10 mL IV as needed for line care before and after medication administration, at discharge, and at de-access ☒ Sodium chloride 0.9% (NS) injection 20 mL IV once as needed for line care post lab draw ☒ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for central line catheter
As Needed Medications	☒ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care ☒ Sodium chloride 0.9% 500 mL continuous infusion at 25 mL/hour IV as needed for line care (TKO for therapy administration)
Referral	☒ Ambulatory referral to OP Infusion Services
PHMC Outpatient Infusion Contact Information	PROVIDER – PLEASE SIGN, DATE AND TIME ORDERS AND RETURN TO: PeaceHealth Peace Harbor Medical Center Outpatient Infusion Services Department 400 Ninth Street, Florence, OR 97439 Contact Phone: 541-902-6019 and FAX 541-902-1649
Authorization by Verbal or Telephone Order	Person giving verbal or telephone order: _____ Person receiving verbal or telephone order: _____ ☐ Check to indicate verbal or telephone orders have been read back to confirm accuracy

Practitioner Signature: _____ **Date of Order:** _____ **Time:** _____

Final page of orders must include signature of the ordering practitioner, date, and time.