



Thank you for selecting our infusion services team to care for your patient. Please provide ALL information listed below to ensure that we can process orders and schedule your patient for treatment without delay.

Part A- Patient scheduling and contact information:

Patient Name (Last, First): _____ Date of Birth: _____

Patient Contact Information and Phone Number(s): _____

Ordering Provider Name (Print): _____

Provider Clinic or Service Address: _____

Clinic or Service Phone Number: _____ Clinic or Service Fax Number: _____

Diagnosis (include ICD 10 codes): _____

Medication and Service Requested - list J-Code/CPT code if known: _____

Date Service is Requested to Begin: _____ **Date Service is Expected to End:** _____

Order will expire 1 year from date of provider signature unless "date service is expected to end" is earlier.

Part B- Insurance and Prior Authorization. Any non-PeaceHealth provider must obtain prior authorization prior to service. Attach a copy of authorization documentation received from insurance payer when submitting orders.

Insurance (Payer) Company: _____

Prior Authorization Number and Conditions: _____

Prior Authorization Expiration Date: _____

Insurance (Payer) Contact Phone Number: _____

Part C- Elements needed to guide medication therapy are included with request for service:

- ☐ Orders and instruction (use the PeaceHealth approved ordering form if you are not a PeaceHealth provider) are complete and include provider signature at the bottom of each page. Check the boxes of ALL orders you would like to activate.
- ☐ For blood products, PeaceHealth Blood and Transfusion Consent form is signed and dated by the provider and the patient.

If information is located outside of PeaceHealth's electronic medical record system attach the following:

- ☐ A list of current medications reconciled by patient provider is available and includes a list of known allergies.
- ☐ Recent progress notes from ordering provider.
- ☐ A copy of relevant laboratory results and other appropriate supporting documentation.

IMPORTANT MESSAGE TO PROVIDERS: To reduce delays in treatment and added phone calls, you may participate to utilize PeaceHealth preferred medication formulary options by signing this document. A clinical pharmacist will adjust orders to align with PeaceHealth medical staff approved medication formulary options, policies, and procedures.

I agree to utilize PeaceHealth preferred medication formulary options, policies and procedures that have been authorized by PeaceHealth Medical Staff. This agreement will be issued for the duration of active orders contained within this treatment plan.

PROVIDER SIGNATURE: _____ **DATE:** _____ **TIME:** _____

FAX completed service request and orders to: PHMC OP Infusion and Nursing Services 541-902-1649



Golimumab (Simponi Aria) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content
For Admission to Service	Provider Instruction – Review information below and address requirements for admission to service: 1. Provider has verified that patient is up to date with all immunizations prior to initiation. 2. Provider has screened patient for tuberculosis, hepatitis, chronic infection, and malignancy prior to initiation. Recommend PPD or Quantiferon TB along with Hep B done within last 1-2 years. 3. If patient on Methotrexate or Leflunomide, then order the following labs: CBC with auto differential and CMP.
Labs	☐ CBC with automated differential once prior to beginning treatment and every 8 weeks ☐ Comprehensive metabolic panel once prior to beginning treatment and every 8 weeks ☒ Treatment lab instructions – Provider approves to release and draw labs 2 days pre and post this planned treatment date.
Pre-Medications	☐ Acetaminophen (Tylenol) tablet 650 mg PO once prior to infusion ☐ Diphenhydramine (Benadryl) 25 mg PO once prior to infusion, OR ☐ Diphenhydramine (Benadryl) 25 mg IV once prior to infusion
Supportive Care	☒ Golimumab (Simponi Aria) 2 mg/kg IV over 30 minutes Select Frequency: ☐ Initiation regimen administered at 0 and 4 weeks followed by maintenance infusion every 8 weeks ☐ Maintenance infusion every 8 weeks Additional order instruction: ☒ Use an infusion set with an in-line low protein-binding 0.22 micron filter. Do not infuse in the same line with other medications.
Nursing Orders	☒ Patients should be evaluated for tuberculosis risk factors and latent tuberculosis infection (with a tuberculin skin test) prior to and during therapy.
Nursing IV Access and Maintenance	Select the most appropriate option below: ☒ Insert PERIPHERAL IV as needed ☒ Sodium chloride 0.9% (NS) flush 10 mL IV once as needed for line care ☐ Access and use NON-PICC Central Line/CVAD as needed and confirm patency ☒ Initiate Central Line (Non-PICC) maintenance protocol ☒ Sodium chloride 0.9% (NS) injection 10 mL IV as needed for line care before and after medication administration, at discharge, and at de-access ☒ Sodium chloride 0.9% (NS) injection 20 mL IV once as needed for line care post lab draw ☒ Heparin, porcine (PF) 100 unit/mL flush 5 mL IV as needed for line care, for de-access ☒ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for clearing central line catheter. Reconstitute with 2.2 mL sterile water for injection to the vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1mg/mL. Instill medication in non-functional lumen. Do not use lumen while dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time not to exceed 120 minutes. Use second dose of Alteplase (Cathflo) if catheter not patent after 120 minutes. If the catheter is functional, aspirate and waste the medication and residual clot prior to flushing line.

Practitioner Signature: _____ **Date of Order:** _____ **Time:** _____

Final page of orders must include signature of the ordering practitioner, date, and time.



Golimumab (Simponi Aria) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content
	☐ Access and use <u>PICC Central Line/CVAD</u> as needed and confirm patency ☒ Initiate PICC maintenance protocol ☒ Change PICC line dressing weekly and as needed ☒ Sodium chloride 0.9% (NS) injection 10 mL IV as needed for line care before and after medication administration, at discharge, and at de-access ☒ Sodium chloride 0.9% (NS) injection 20 mL IV once as needed for line care post lab draw ☒ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for clearing central line catheter
As Needed Medications	☒ Acetaminophen (Tylenol) tablet 650 mg PO once as needed for aches or temp change > 2 degrees F ☒ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care ☒ Sodium chloride 0.9% 500 mL continuous IV infusion at 25 mL/hour as needed for line care (TKO for therapy administration)
Emergency Medications	If patient has symptoms of anaphylaxis (wheezing, dyspnea, hypotension, angioedema, chest pain, or tongue swelling), discontinue infusion and initiate standard emergency response procedures. ☒ DiphenhydrAMINE (Benadryl) injection 25-50 mg IV once as needed for mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (greater than or equal to 20 points in SBP), nausea, urticaria, chills, pruritis). - Administer 50 mg IV if patient has NOT had diphenhydramine within 2 hours of reaction - Administer 25 mg IV if patient has had diphenhydramine within 2 hours of reaction, if reaction doesn't resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and notify provider. ☐ Albuterol 90 mcg/actuation inhaler 2 puffs once as needed for wheezing, shortness of breath associated with infusion reaction and notify provider. Administer with a spacer if available. ☒ MethylPREDNISolone (Solu-Medrol) injection 125 mg IV once as needed for shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (greater than or equal to 20 points in SBP), nausea, urticaria, chills, pruritis) that worsen or persist after administration of diphenhydramine (Benadryl) and notify provider. ☒ EPINEPHrine (Adrenalin) injection 0.5 mg IM once as needed for severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (greater than or equal to 40 points in SBP), shortness of breath with wheezing and O2 Sat less than 90%), and notify provider.
Referral	☒ Ambulatory referral to OP Infusion Services
PHMC Outpatient Infusion Contact Information	PROVIDER – PLEASE SIGN, DATE AND TIME ORDERS AND RETURN TO: PeaceHealth Peace Harbor Medical Center Outpatient Infusion Services Department 400 Ninth Street, Florence, OR 97439 Contact Phone: 541-902-6019 and FAX 541-902-1649
Authorization by Verbal or Telephone Order	Person giving verbal or telephone order: _____ Person receiving verbal or telephone order: _____ ☐ Check to indicate verbal or telephone orders have been read back to confirm accuracy

Practitioner Signature: _____ **Date of Order:** _____ **Time:** _____

Final page of orders must include signature of the ordering practitioner, date, and time.